

Oral Health Integration in Pediatric Primary Care and the Age One Dental Visit: A Mixed Method Study



THE OHIO STATE UNIVERSITY
COLLEGE OF DENTISTRY



Natalie Bussard, DDS^{1,2}; Paul Casamassimo, DDS, MS^{1,2}; Beau Meyer, DDS, MPH^{1,2}; Andrew Wapner, MD, MPH^{1,3}; Jin Peng, MD, MS, PhD¹; Homa Amini, DDS, MS, MPH^{1,2}
1.The Ohio State University College of Dentistry 2. Nationwide Children's Hospital 3. The Ohio State University College of Public Health

PURPOSE

- Compare age of first dental visit between children who did and did not receive preventive oral health services at their primary medical home
- Understand pediatrician's attitudes, beliefs, and barriers to the implementation of oral health integration

METHODS

Quantitative

- Retrospective claims data from Medicaid-enrolled children 1-5 years of age in Ohio
- Exposure: physician-applied fluoride varnish (PAFV)
- 1° outcome: age of first dental visit
- 2° outcomes: subsequent dental utilization
- T-test

Qualitative

- 10 semi-structured interviews with pediatricians in rural and metro Ohio
- Inductive thematic analysis

RESULTS

Receipt of FV by physician	N	Mean Age, y (95% CI)	Pr > t
No	24342	4.4 (4.3, 4.4)	<.0001
Yes	8951	3.5 (3.5, 3.6)	

- PAFV group had lower frequency of caries-related dental treatment visits (39% vs. 59%)

Oral health integration in pediatric primary care is associated with dental visits at a younger age.

Pediatricians desire oral health provider integration, but significant barriers exist.





Figure 1. Thematic summary of qualitative interviews regarding barriers and drivers of oral health integration into primary care

Scan to access interview guide and references.



Acknowledgements

- Susan Lawson with Ohio Assoc. of Community Health Centers
- Charitha Gowda, MD, MPH; acting medical director of PFK at time of study design
- Data used in this study was partially provided by Partners For Kids, a pediatric accountable care organization established by Nationwide Children's Hospital.

Disclaimer

The statement in this study are solely the responsibility of the authors and do not necessarily represent the views of Partners For Kids or Nationwide Children's Hospital.

Funding

This project was completed without specific funding.